

Cytotoxic and Angiosuppressive Potentials of *Zehneria japonica* (Thunb. ex Murray) S.K. Chen (Cucurbitaceae) Crude Leaf Extracts

Marri Jmelou Roldan^{1,4*}, Ting-Yu Chin⁵, Yun-Chieh Tsai⁵, Agnes L. Castillo^{1,2,4}, Oliver B. Villaflores^{1,3,4}

*Corresponding author's email address: mjmeroldan@gmail.com

¹ The Graduate School, University of Santo Tomas, Manila, Philippines,

² Faculty of Pharmacy, University of Santo Tomas, Manila, Philippines,

³ Department of Chemistry, College of Science University of Santo Tomas, Manila, Philippines,

⁴ Research Center for the Natural and Applied Sciences, University of Santo Tomas, Manila, Philippines,

⁵ Department of Biotechnology, Chung Yuan Christian University, Chungli District, Taoyuan City, Taiwan

RESEARCH ARTICLE

Abstract

Background and Objectives: *Zehneria japonica* belongs to the Cucurbitaceae family which is one of the most important plant families. It is commonly known as "Pipinong-gubat," widely distributed in Central Luzon regions and in areas along streams and clearings at low and medium altitudes in the Philippines. This study aimed to evaluate the potential cytotoxic and angiosuppressive properties of *Zehneria japonica* (Thunb. ex Murray) S.K. Chen (Cucurbitaceae) leaf extracts.

Methodology: The *Z. japonica* semi-crude extracts were obtained by sequential extraction using hexane, ethyl acetate, and n-butanol. A modified duck egg chorioallantoic membrane (CAM) assay was aided by AngioQuant, a digital imaging software used to evaluate angiogenic activity. Inhibition of angiogenesis was evaluated by percent increase or decrease in mean length of blood vessels, mean size of blood vessels, and total number of blood vessel junctions. Moreover, the cytotoxic effects of the extracts were determined through MTT Assay. Osteosarcoma (U2Os) and hepatocellular carcinoma (HepG2) cells were used as cancer representatives while human umbilical vein endothelial cells (HUVEC) were used as the normal cell control.

Results: Analysis with AngioQuant revealed that treatment of the duck egg CAM with *Z. japonica* semi-crude extracts suppressed angiogenesis with IC₅₀ values of 1,810.00 µg/mL, 192.50 µg/mL, and 147.70 µg/mL for hexane, ethyl acetate, and n-butanol, respectively, with Celecoxib (20 µg/mL) as the positive control. For MTT assay, *Z. japonica* extracts exhibited strong cytotoxic effect against U2Os with an IC₅₀ values of 19.65 µg/mL, 9.89 µg/mL, and 31.04 µg/mL for the hexane, ethyl acetate, and n-butanol extracts, and no cytotoxic effects against HepG2 with IC₅₀ values of 770.90 µg/mL, 130.10 µg/mL and 231.60 µg/mL for the hexane, ethyl acetate, and n-butanol extracts. Doxorubicin (0.544 µg/mL) was used as the positive control. The extracts also inhibited the growth of the normal cells, with IC₅₀ values of 69.46 µg/mL, 42.23 µg/mL and 63.44 µg/mL for the hexane, ethyl acetate, and n-butanol extracts. There were no mortality and toxic symptoms observed for 14 days after the administration of the crude butanolic extract of *Z. japonica* in six female Sprague-Dawley rats.

Conclusion: *Z. japonica* crude leaf extracts exhibited angio-suppressive activity through CAM assay. In MTT assay, the extracts exhibited strong cytotoxicity in U2Os (IC₅₀ ≤ 20 µg/mL), no cytotoxic effect in HepG2 (IC₅₀ > 100 µg/mL) cells, and mild cytotoxic effect in HUVEC (IC₅₀ 40-60 µg/mL). Phytochemical screening through TLC revealed that the extracts contain alkaloids, anthrones, flavonoids, and sterols.

Keywords: *Zehneria japonica* (Thunb. ex Murray) S.K. Chen, Cucurbitaceae, cytotoxic, MTT Assay, angiosuppressive, CAM Assay

Introduction

The vast increase in new cases of various diseases has been detrimental to one's health and lessens the person's quality of life. One of the notable diseases that causes morbidity and mortality worldwide is cancer. According to the World Cancer Report, there were approximately of 8.2 million cancer-related deaths and 14 million new cases of cancer in

2012 [1]. Cancer cells grow abnormally and they frequently invade healthy cells in the body [2]. Angiogenesis plays a vital role in the etiology of tumor progression and metastasis [3]. Innovation of new strategies and therapies in the field of oncology has led to the study of angiogenic inhibitors [4]. Synthetic angiogenic inhibitors have been produced like Bevacizumab but investigations involving natural sources or products have limited or few studies available [5].

Through the years, plants have been utilized as sources of useful antitumor compounds that have immensely contributed in the fight and prevention against cancer which have proved to be promising. The Philippines has a diverse collection of plants but these resources are not yet well explored in pursuit of novel antiangiogenic agents [6]. A keen interest in discovering an alternative source of anti-cancer agents from medicinal plants had led to the study of the Cucurbitaceae family.

Zehneria is one of 125 genera in the family Cucurbitaceae. *Zehneria* species have been utilized in traditional medicine to treat fever, stomach pain, diarrhea, skin diseases, jaundice, and kidney dysfunction [7,8]. Secondary metabolites reported from this genus include dammarane-type glycosides (gypenosides) and sphingolipids with antimicrobial and antitumor activities [8,9].

Zehneria japonica (Thunb. ex Murray) S.K. Chen is widely distributed in Central Luzon region of the Philippines where it is locally known as "Pipinong-gubat." No study has been made to explore its anti-angiogenic and cytotoxic properties although other plants in the same family were proven to possess the anticipated pharmacological activities.

This study aimed to evaluate the cytotoxic and angiosuppressive property of *Z. japonica* which may be a potential source of phytochemical active compounds with the use of the duck chorioallantoic membrane (CAM) and MTT colorimetric assay, respectively.

Methodology

Instrumentation

Spectrophotometric measurements were performed by using Shimadzu RF-1501 UVVIS (ultraviolet-visible spectrophotometer). Cytotoxic activity of the crude extracts was determined using MTT assay and was measured spectrophotometrically (Biotek® ELX808 Absorbance Microplate Reader). Inverted microscope (Olympus CKX41 microscope), centrifuge (Eppendorf 5810R), and CO² incubator (5% CO², 95% humidity and 37°C; Thermo Scientific TM 8000 incubator) were also used. INCUBOX™ automatic humidified incubator was used for CAM assay.

Chemicals

Technical grade reagents bought from Belman laboratories were used for extraction. Analytical grade reagents were used for the phytochemical screening and biological assays of the air-

dried, ground leaves of *Z. japonica*. Celecoxib was used for the positive control group while dimethylsulfoxide (DMSO) was used for the negative group in the CAM assay.

Plant Material

Fresh leaves of *Zehneria japonica* were collected in March 2016 at Basista, Pangasinan. The plant sample was identified and authenticated by Asst. Prof. Ophelia Laurente, curator at the University of Santo Tomas Herbarium, where a voucher specimen (No. USTH- 013124) has been deposited. The leaves were air-dried under shade for three weeks until they became brittle. The leaves were pulverized in a Wiley mill grinder and weighed.

Test animals

A total of six female, non-pregnant, nulliparous Sprague-Dawley rats, aged 8–12 weeks, weighing 100–160g were selected for the toxicity test. The test animals were procured from the Food and Drug Administration in Alabang, Muntinlupa City, Philippines, and were kept at an animal house facility, Research Center of Natural and Applied Sciences, University of Santo Tomas. The animals were subjected to a 7-day acclimatization. An approval from the Institutional Animal Care and Use Committee was obtained as a requirement for the animal testing.

Egg Preparation

Fertile duck eggs were obtained from an egg farm in Baliuag, Bulacan, Philippines. A total of twelve groups composed of five zero-day old eggs were used in the assay. The eggs were incubated with the use of the Incubox automatic turning model with a one hour rolling interval at (38 ± 1) °C in a humidified incubator until the Embryonic Development Day (EDD). The eggs were opened on either 8th, 9th, or 10th day of incubation. Tap water was placed beneath the egg trays which had a wire mesh bottom to prevent excessive humidity that might allow surface organisms to penetrate the shell [10, 11].

Cell Lines, Culture Treatments

Osteosarcoma (U2Os), Hepatocellular carcinoma (HepG2), and HUVEC lines cultured by recommended protocols of ATCC were obtained from the Department of Biotechnology in Chung Yuan Christian University, Taoyuan, Taiwan and were applied to the plant hexane, ethyl acetate, and n-butanol extracts to determine their cytotoxic activity by the MTT colorimetric assay. The cells were seeded at 104 cells/100 µL into each well of 96-well plates and incubated for 24 hours at 37°C and in 5% CO². The culture medium was removed and the extracts were added to

the wells in different concentrations (2, 10, 20, 100, 500 µg/mL and positive control Doxorubicin 0.544 µg/mL).

Preparation of Plant Extracts

A total of 1.17 kg of the ground leaves was obtained. The powder was extracted sequentially using solvents of increasing polarities. One kilogram of powdered leaves was soaked in 4L hexane (non-polar solvent) for 24 hours at room temperature. After one-day, the hexane extract was collected and the procedure was repeated thrice. The marc/residue was dried before the addition of the next solvent. Then, subsequent extractions with ethyl acetate (semi-polar solvent) and n-butanol (polar solvent) were done in the same sample. The three semi-crude extracts were concentrated using a rotary evaporator and water bath (40°C) to remove the excess solvent. The extracts were properly labeled and stored at 4°C for further use.

Phytochemical Screening

Prepared extracts were subjected to preliminary phytochemical screening in order to qualitatively determine some of the secondary metabolites: saponins, terpenoids, flavonoids, tannins, alkaloids, quinones, anthraquinones, and reducing sugar using appropriate methods [12,13,14].

Phytochemical screening was done by using a thin layer of chromatography sprayed with color reaction reagents. The sample was applied using a capillary tube on a strip of pre-coated silica plates (Merck EMD TLC). After the sheet was air-dried, it was developed in a chamber that was a beaker containing the most suitable solvent covered with a watch glass. For the *Z.japonica* Hexane (ZjH), a solvent system of hexane and ethyl acetate (7:3) was used. On the other hand, *Z. japonica* Ethyl acetate (ZjEA) was developed with hexane and ethyl acetate (5:5), while *Z.japonica* n-Butanol (ZjB) was developed with dichloromethane and methanol (9:1). The developed chromatogram was air-dried and visualized by spraying with a suitable reagent for the desired constituents.

Toxicity Test

The ZjB was assessed for toxicity by using the acute oral toxicity test according to the Organization for Economic Cooperation and Development Guideline 425 [10]. One dose level of 2,000 mg/kg body weight was performed. Initially, the animals were selected randomly and equally divided into two groups designated as Group A (1 rat) and Group B (4 rats). The rats were marked for proper identification and

were subjected to a 7-day acclimatization. Prior to dosing, the rats were fasted overnight (12 hours) by withholding food but not water. The fasted weight was recorded and the suitable dose for each rat was computed. The ZjB to be administered was diluted with water. One dose level of 2,000 mg/kg body weight was given to Group A via oral gavage. General toxicity signs and behavioral changes were observed daily for a total of 14 days and were noted. With no mortality in a five-day observation period, Group B was given a one dose level of 2,000 mg/kg body weight. The same observation done on Group A was executed for Group B. After the testing period, the rats were sacrificed through asphyxia using carbon dioxide and were subjected to gross necropsy with the assistance of a Veterinarian, Dr. Joy Cabigan.

Chorioallantoic Membrane Assay

Chorioallantoic Membrane assay was used to determine the angiosuppressive activity of the plant extracts. It was performed based on the method of West *et al.* [11,15] with minor modifications on the choice of control agents. Celecoxib was used as a positive control agent and DMSO as vehicle control.

On the 8th day of incubation, the surface of the egg was cleaned with 70% alcohol. A small window was made at the side of the egg using a drill bit and forceps. The filter discs with the test solution (10, 100, and 1000 µg/mL of hexane, ethyl acetate, and n-butanol crude extracts and the controls: DMSO and Celecoxib 20 µg/mL) were placed on top of the CAM and the window was sealed using a clear tape. Afterwards, the egg was incubated for another 48 hours.[16,17]. After incubation, the egg was opened with the general condition of the embryo

$$\% \text{ inhibition} = \frac{\text{no. of branch points (treated)} - \text{no. of branch points (control)}}{\text{no. of branch points (control)}} \times 100$$

and CAM photographed and quantified using the software, AngioQuant™ [11]. The same procedure was applied for the rest of the eggs. The percent inhibition was determined in each test group using the formula:

MTT Cytotoxicity Assay

MTT assay was performed to determine the cell viability which was measured by the reduction of MTT to a purple colored formazan product [18,19]. After 24 hours of treatment, 20 µL MTT reagent was added to each well. After incubation for four hours at 37°C in the dark, the supernatants

of culture media was aspirated and 150 μ L DMSO was added to each well to dissolve the formazan crystals. Then, the absorbance was taken at 590nm using a microplate reader (Biotek® ELX808).

Statistical method

The means and its standard error were used to summarize the percentage inhibition of hexane, ethyl acetate, and n-butanol at different concentrations in CAM and MTT Assay. Single-factor analysis of variance was used to determine the effect of different concentrations of each extract in the percentage inhibition with Tukey's HSD for Post Hoc analysis (MTT). Celecoxib was further compared with each of the concentrations (CAM). Meanwhile, estimates of IC_{50} were determined using four parameter logistic regression model. All these tests were performed using SPSS ver. 20.0 and GraphPad Prism 6.0. All tests were set at 5% level of significance.

Results and Discussion

This study determined and evaluated the cytotoxic and angiosuppressive potentials of *Zehneria japonica* (Thunb. ex Murray) S.K. Chen hexane, ethyl acetate, and n-butanol crude leaf extracts.

Plant Extraction

The dried leaves of *Z. japonica* weighing 1.17 kg obtained a percentage yield of 1.68% for hexane crude extract, 2.56% for ethyl acetate crude extract, and 0.53% for n-butanol crude extract.

Phytochemical Screening

In the preliminary phytochemical screening using thin layer chromatography, the leaves of *Z. japonica* crude extracts revealed the presence of various secondary metabolites (Table 1). Among those detected in the crude extracts were alkaloids, anthrones, flavonoids, and phenols.

Acute Oral Toxicity Test

The animals showed no mortality and no change in behavior for 14 days after the administration of the crude butanolic extract of *Z. japonica*. Gross necropsy showed no significant findings observed on all limit test animals upon opening and examining its body cavities (Figure 1). Results obtained from the acute oral toxicity test showed that *Z. japonica* was practically

Table 1. Phytochemical screening of *Z. japonica* crude extracts

	ZjH	ZjEA	ZjB
Alkaloids	+	++	+
Anthrones	++	+	++
Flavonoids	++	++	+
Phenols, triterpenes, sterols	+	++	++
Sugar	-	+	+

LEGEND: ++: strong positive test; +: low positive test; -: negative test




Figure 1. Gross Necropsy of the female Sprague-Dawley rats for Acute Oral Toxicity

non-toxic at a dose of 2,000mg/kg BW.

Chorioallantoic Membrane Assay

The difference in the vascularization of the egg treated with different semi-crude extracts of the plant was assessed at a macroscopic level in Figure 2. Representative chorioallantoic membrane of the samples incubated for 10 days at 38°C +/- 1°C of a humidified incubator with 1,000 µg/disc concentration were chosen because at this dose level, the highest anti-angiogenic response was observed. There was an evident reduction in the number of blood vessels as compared with the negative control group, which indicates that the growth of new blood vessels had been prevented.

The means of groups were compared and analyzed using single factor analysis of variance (ANOVA). The treatment groups revealed different responses based on the mean blood vessel count as presented in Figure 3. The p-values of all the extracts was >0.05 thus, considered as statistically insignificant (Table 2). This means that the three extracts have comparable angiosuppressive properties at 1000, 100, and 10 µg/mL respectively.

The data obtained from the analysis showed that all the extracts exhibited a response that is comparable to the positive control, Celecoxib. Thus, the ZjH, ZjEA, and ZjB in all the given concentrations showed anti-angiogenic effect. The graph in Figure 4 reveals a dose-dependent activity for the percent inhibition of the three (3) extracts. This indicates a direct proportional relationship that implies that as the dose of a drug increases, the activity also increases. In the same





Figure 2. Representative chorioallantoic membrane of the samples incubated for 10 days at 38°C +/- 1°C of a humidified incubator with 1000 µg/disc concentration. Sprouting of secondary blood vessels lessened and become thinner in the 1000 µg/disc of the extracts and in the positive control, Celecoxib when compared to DMSO. The resolution of the camera has a great impact in quantifying the blood vessels when using the AngioQuant™ to determine the Mean ± SD blood vessels.



Legend: CELE - Celecoxib; ZjH - *Z. japonica* Hexane Extract; ZjE - *Z. japonica* Ethyl acetate Extract; ZjB - *Z. japonica* Butanol Extract

Figure 3. The anti-angiogenic activity of *Z. japonica* extracts represented by mean blood vessel count using AngioQuant™


Table 2. Percentage Inhibition of the *Z. japonica* extracts at different concentrations in CAM assay.

Concentration					
	10 ug/mL	100 ug/mL	1000 ug/mL	F stat	p-value
Hexane	27.53±4.29	36.46±8.50	47.32±7.88	1.929	0.201
Ethyl acetate	35.87±12.39	47.99± 12.29	57.14±10.85	0.808	0.476
Butanol	36.25±13.72	46.26±11.63	61.34±12.24	1.011	0.402

There is no significant difference in the mean percentage inhibition of hexane [F2,9 = 1.929, p=0.201], ethyl acetate [F2,9 = 0.808, p=0.476], and butanol [F2,9 = 1.011, p=0.402] at different concentrations. Moreover, all the mean percentage inhibitions of hexane [F2,9=1.929, p = 0.201], ethyl acetate [F2,9 = 0.808, p = 0.476], and butanol [F2,9=1.011, p = 0.402] at different concentrations did not differ with Celecoxib. Values expressed as mean ± SEM, n = 4.

Table 3. The IC_{50} values of the different extracts based on the four-parameter logistic regression model.

Estimate			
	IC_{50}	95% Confidence Interval	R^2
Hexane	1810.0	156.7 – 20,907	0.922
Ethyl acetate	192.5	18.5 – 2,008	0.805
Butanol	147.7	19.0 – 1,148	0.789



Legend: CELE - Celecoxib; ZJH - *Z. japonica* Hexane Extract; ZJE - *Z. japonica* Ethyl acetate Extract; ZJB - *Z. japonica* Butanol Extract


Figure 4. The effect of *Z. japonica* extracts on angiogenesis “expressed as mean percent inhibition”.

figure, it has revealed that ZJB at 1,000 µg/mL/disc yielded the highest activity with 61.34% inhibition which is comparable with Celecoxib at 20 µg/mL/disc, with a 61.90% inhibition. At 1,000 µg/mL/disc concentration of all the extracts, ZJH indicated the least activity with 47.32% inhibition.

The IC_{50} estimates for the extracts were obtained using the four-parameter logistic regression model as shown in Table 3. Among the different extracts, hexane is the least potent with 1,810 µg/mL as compared to ethyl acetate (192.5 µg/mL) and n-butanol (147.7 µg/mL) which is the most potent.


MTT Cytotoxicity Assay

The 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyl tetrazolium bromide (MTT) assay is based on the principle that only viable and metabolically active cells can cleave the MTT dye to produce formazan crystals, which is dissolved and quantified using a spectrophotometer (20). Figure 5 shows the cytotoxic effect of *Z. japonica* extracts expressed as mean percent inhibition. Among the three extracts, *Z. japonica* hexane extract in 100 µg/mL exhibited the highest inhibition with 98.41% followed by ethyl




Legend: DOXO - Doxorubicin; ZJH - *Z. japonica* Hexane Extract; ZJE - *Z. japonica* Ethyl acetate Extract; ZJB - *Z. japonica* Butanol Extract

Figure 5. Cytotoxic potential of *Z. japonica* crude leaf extracts against U20s cells expressed as Mean percent inhibition.



Legend: DOXO - Doxorubicin; ZJH - *Z. japonica* Hexane Extract; ZJE - *Z. japonica* Ethyl acetate Extract; ZJB - *Z. japonica* Butanol Extract

Figure 6. Cytotoxic potential of *Z. japonica* crude leaf extracts against HepG2 cells expressed as Mean percent inhibition.



Legend: DOXO - Doxorubicin; ZJH - *Z. japonica* Hexane Extract; ZJE - *Z. japonica* Ethyl acetate Extract; ZJB - *Z. japonica* Butanol Extract

Figure 7. Cytotoxic potential of *Z. japonica* crude leaf extracts against HUVECs cells expressed as Mean percent inhibition.

acetate extract in 100 µg/mL with 96.62%. In HepG2, the extracts exhibited a dose-dependent activity and ethyl acetate extract showed the highest inhibition with 77.75% much higher than the positive control and the two other extracts (Figure 6). And lastly, in HUVEC (normal cells), n-butanol extract exhibited the highest inhibition of 108.08% among the three extracts (Figure 7).

Based on the statistical treatment, the mean percentage of the three crude extracts exhibited no significant difference (p-values <0.05) as compared with the positive control, Doxorubicin. However, a comparison of the mean percentage of the three extracts in different concentrations showed a significant difference. For U2Os cells, the mean percentage of 2 µg/mL was significantly less (p-value <0.011) than 20 µg/mL, and that the mean percentage of the 100 µg/mL was significantly higher (p-value <0.001) than 20 µg/mL. For HepG2 cells, it was evident that the mean percentage of the three extracts significantly differed (p-value 0.037). The mean percentage inhibition of

Doxorubicin, n-butanol, and ethyl acetate extracts did not differ (p-value 0.430), and that the mean percentage inhibition of hexane was significantly the least (p-value <0.05). Additionally, for hexane, ethyl acetate and n-butanol extracts, the mean percentage of the different concentrations significantly differed (p-value <0.001), indicating that the mean percentage of 10 µg/mL is significantly less (p-value 0.011) than the 100 µg/mL, and that the mean percentage of the 500 µg/mL concentration was significantly higher (p-value <0.001) than the 100 µg/mL. Finally, for the HUVEC, there were no significant differences in the mean percentage inhibition of the three extracts and Doxorubicin. Yet, the mean percentage of the concentrations of the three extracts significantly differed (p-value <0.001), indicating that the mean percentage of 10 µg/mL is significantly less (p-value <0.001) than the 100 µg/mL and 500 µg/mL. The mean percentage inhibition of 100 µg/mL and 500 µg/mL did not differ (p-value 0.730).

Table 4. In vitro cytotoxicity effects of *Zehneria japonica* extracts against U2Os, HepG2, and HUVEC.

Test Extracts	ZJH			ZJE			ZJB		
Cells	U2Os	HepG2	HUVEC	U2Os	HepG2	HUVEC	U2Os	HepG2	HUVEC
IC ₅₀ (µg/mL)	19.65	770.90	69.46	9.89	130.10	42.23	31.04	231.60	63.44
R ²	0.813	0.867	0.560	0.825	0.886	0.441	0.653	0.884	0.731

The IC₅₀ values of ZjH, ZjEA, and ZjB for U2Os cells were 19.65 µg/mL, 9.89 µg/mL, and 31.04 µg/mL respectively. Similarly, the IC₅₀ values of ZjH, ZjEA, and ZjB for HepG2 cells were 770.90 µg/mL, 130.10 µg/mL, and 231.60 µg/mL respectively. Lastly, the IC₅₀ values of ZjH, ZjEA, and ZjB for HUVEC were 69.46 µg/mL, 42.23 µg/mL, and 63.44 µg/mL respectively (Table 4).

In summary, the plant *Zehneria japonica* also possesses an anti-angiogenic activity that may be related to other studies of its related plant species under Cucurbitaceae [21,22]. The anti-angiogenic effect of *Z. japonica* may be attributed to phytochemical constituents that are prominent in the three extracts such as sterols, triterpenes, anthrones, flavonoids, and alkaloids.

The polyphenols exhibit anti-angiogenic effect and metastasis through the regulation of multiple signaling pathways. It may also influence some steps in tumor angiogenesis. Similarly, the presence of triterpenoids and sterols have strong anticancer activity [23]. Among the three extracts, n-butanol extract exhibited the highest potency and inhibition in CAM assay comparable to the positive control, Celecoxib.

Cytotoxic compounds directly targeted cells and are utilized in chemotherapy in cancer patients. According to the United States National Cancer Institute, a sample or compound is considered cytotoxic if its IC₅₀ value is ≤ 20 µg/mL. Based on the results, *Z. japonica* extracts exhibited strong cytotoxicity against U2Os cells with an IC₅₀ values within 20 µg/mL except for n-butanol extract with 31.04 µg/mL, and no cytotoxic effect against HepG2 cells with an IC₅₀ values of >100 µg/mL. The extracts exhibited mild cytotoxicity against HUVEC with IC₅₀ values of 40-60 µg/mL. Among the three extracts, ethyl acetate extract has the highest IC₅₀ values based on the three cell lines and is considered the most potent among the extracts. Since it exhibits activity against cancer and normal cells, n-butanol extract is much safer to use than ethyl acetate because its IC₅₀ values were revealed to be in the middle when compared to hexane and ethyl acetate extracts.

The presence of terpenoids and flavonoids is attributed to the cytotoxic activity of *Z. japonica*. Flavonoids, specifically, are reported to have cytotoxic activity due to the presence of phenolic groups, while terpenoids prevent the cancer cell growth and size reduction [24,25].

Conclusion

The results suggest that *Z. japonica* leaf extracts have a cytotoxic and angiosuppressive effect. Phytochemical screening through TLC revealed that the extracts contain alkaloids, anthrones, flavonoids, and sterols. Further studies should evaluate the *in vitro* and *in vivo* benefits of *Z. japonica* extracts.

Acknowledgements

This study was supported by Mr. Will Kian Go of Daido Impex Corporation. The authors are grateful to Prof. Hsi-Ya Huang and Dr. Stephen Lirio of the Department of Chemistry, together with Mr. Meng-Chi Chung and Ms. Hsiang-Chi Chen of the Department of Biotechnology in Chung Yuan Christian University, and the University of Santo Tomas Graduate School and Research Center for the Applied and Natural Sciences.

References

1. World Cancer Report. (2012) Cancer Statistics.
2. De Mesquita ML, De Paula JE, Pessoa C, De Moraes MO, Costa-Lotufo LV, Grougnet R, Michel S, Tillequin F, Espindola LS. (2009) Cytotoxic activity of Brazilian Cerrado plants used in traditional medicine against cancer cell lines. *Journal of Ethnopharmacology* 123:439-45.
3. Nassar ZD, Aisha AF, Ahamed MB, Ismael Z, Au-Salah KM, Alrokayan SA, *et al.* (2011) Antiangiogenic properties of koetjapic acid, a natural triterpene isolated from *Sandoricum koetjaoe* Merr. *Cancer Cell International* 11: 12.
4. Bisacchi D, Benelli R, Vanzetto C, Ferrari N, Tosetti F, Albini A. (2003) Anti-angiogenesis and angioprevention: mechanisms, problems and perspectives. *Cancer Detection and Prevention* 27: 229-38.
5. Herrera AA, Amor EC. (2011) Antiangiogenic activity of extracts and fractions of an endemic plant *Ardisia pyramidalis* (Cav.) Pers. from Bataan, Philippines using duck in ovo chorioallantoic membrane assay. *Journal of Medicinal Plants Research* 5(13): 2637-46
6. Tantiado RG, Tan VP. (2012) Evaluation of the angiosuppressive activity of *Tinospora rumphii* Boerl. Stem extract using the chorioallantoic membrane assay in *Anas platyrhynchos* embryos. *International Journal of BioSciences and BioTechnology* 4(2): 93-100.
7. Kirtikar Basu BD. (1975) Indian medicinal plants,

- vol. II. New Delhi: Jayyed Press; p. 1161-62.
8. Anand SP, Jeyachandran R, Nandagopalan V. (2011) NMR spectral analysis on root extract of *Zehneria scabra* a vital medicinal climber. *Journal of Pharmaceutical Sciences and Research* 3:1015-18.
9. Li H, Luo Y, He Z, Zhang G. (2006) Phytochemical study on *Zehneria maysorensis*. *Natural Product Research and Development* 18:411-14.
10. Organization for Economic Co-operation and Development. (2001) OECD guidelines for the testing of chemicals. Acute oral toxicity – acute toxic class method. Paris: Organization for Economic Co-operation and Development.
11. Reña S. (2016) The anti-angiogenic and antibacterial effect of *Tinomisium philippinense* Miers. (Menispermaceae) leaf extract. *Journal of Coastal Life Medicine* 4(1):61-68.
12. Benariba N, Djaziri R, Bellakhdar W, Belkacem N, Kadiata M, Malaisse WJ, *et al.* (2013) Phytochemical screening and free radical scavenging activity of *Citrullus colocynthis* seeds extracts. *Asian Pacific Journal of Tropical Biomedicine* 3(1): 35-40.
13. Benarba B, Meddah B, Aoues A. (2012) *Bryonia dioica* aqueous extract induces apoptosis through mitochondrial intrinsic pathway in BL41 Burkitt's lymphoma cells. *Journal of Ethnopharmacology* 141: 510-6.
14. Khan AM, Qureshi RA, Ullah F, Gilani SA, Nosheen A, Sahreen S, *et al.* (2011) Phytochemical analysis of selected medicinal plants of Margalla Hills and surroundings. *Journal of Medicinal Plants Research* 5(25): 6055-60.
15. West DC, Thompson WD, Sells PG, Burbridge MF. (2001) Angiogenesis assays using chick chorioallantoic membrane. In: Murray JC, editor. *Methods in molecular medicine*. New Jersey: Human Press; p. 107-29.
16. Ribatti D. (2010) The Chick embryo Chorioallantoic membrane in the study of angiogenesis and metastasis. Springer Science & Business Media, Dordrecht.
17. Blacher S, Devy L, Hlushchuk R, Larger E, Lamande N, Burri P, Corvol P, Djonov V, Foidart JM, Noel A. (2005) Quantification of Angiogenesis in the chicken chorioallantoic membrane (CAM). *Image Analysis and Stereology*. 24: 169-180.
18. Sleman K, Mahmoud R, Anwar R. (2013) Anticancer activity of Anise (*Pimpinella anisum* L.) seed extract. *Open Nutraceuticals J.* 6:1-5.
19. Rupachandra S, Sarada DVL. (2013) Anticancer activity of methanol extract of the seeds of *Borreria hispida* and *Momordica dioica*. *Journal of Pharmaceutical Research*, 6:565-568.
20. Stockert JC, Blazquez-Castro A, Cañete M, Horobin RW, Villanueva A. (2012) MTT assay for cell viability: Intracellular localization of the formazan product is in lipid droplets. *Acta Histochemica* 114(8):785-796.
21. Ayyad SN, Abdel- Lateff A, Alarif WM, Patacchioli FR, Badria FA, Ezmirly ST. (2012) In-vitro and in-vivo study of cucurbitacin s-type triterpene glucoside from *Citrullus colocynthis* growing in Saudi Arabia against hepatocellular carcinoma. *Environmental Toxicology and Pharmacology* (33):245-251.
22. Kongue MDT, Talontsi FM, Lamshoft M, Kenla TJN, Dittrich B, Kapche GDWF, Spiteller M. (2013) Sonhafouonic acid, a new cytotoxic and antifungal hopene-triterpenoid from *Zehneria scabra camerunensis*. *Fitoterapia* (85): 176-180.
23. Raturi P, Sharma A. (2011) Natural compounds in the treatment of cancer. *International Journal of Science and Technology and Management* 2:79-85.
24. Bhavani MB, Leelavathi S. (2015) Investigation on in vitro cytotoxic activity of a selected wild Cucurbitaceae plant *Corallocarpus epigeus* against cancer. *International Journal of Pharmaceutical Sciences and Research* 51:3554-57.
25. Sudarsanam D, Helen Mary L, Lakshmi, Tilton F, Joseph J, Siddharta SS. (2014) Phytochemical, antioxidant, and cytotoxic properties of the fruit extract from *Cucurbita digitata*. *International Journal of Pharmacy and Pharmaceutical Sciences* 6(4):353-356.